



# Functional significance of ultraviolet feeding cues in wild turkeys

Scott J. Werner <sup>a,\*</sup>, Richard Buchholz <sup>b</sup>, Shelagh K. Tupper <sup>a</sup>, Susan E. Pettit <sup>a</sup>, Jeremy W. Ellis <sup>a</sup>


<sup>a</sup> USDA/APHIS/WS/National Wildlife Research Center, 4101 LaPorte Avenue, Fort Collins, CO 80521-2154, United States

<sup>b</sup> University of Mississippi, Biology Department, 104 Shoemaker Hall, University, MS 38677-1848, United States

## HIGHLIGHTS

- Wild turkeys do not prefer UV feeding cues regardless of feeding experience.
- UV feeding cues are used functionally for avian foraging behavior.
- Postingestive consequences are necessary for conditioned avoidance of UV feeding cues.
- Intestinal parasite infection influences the process of food selection in wild turkeys.

## GRAPHICAL ABSTRACT



## ARTICLE INFO

### Article history:

Received 4 November 2012

Received in revised form 15 October 2013

Accepted 18 October 2013

### Keywords:

Conditioned avoidance

*Eimeria* spp.

Foraging behavior

*Meleagris gallopavo*

Postingestive consequence

Visual cue

## ABSTRACT

Most birds are able to sense ultraviolet (UV) visual signals. Ultraviolet wavelengths are used for plumage signaling and sexual selection among birds. The aim of our study was to determine if UV cues are also used for the process of food selection in wild turkeys (*Meleagris gallopavo*). We used avoidance conditioning to test the hypothesis that UV feeding cues can be used functionally for foraging behavior in wild turkeys. Female turkeys exhibited no avoidance of untreated food and 75–98% avoidance of food treated with an UV-absorbent, postingestive repellent (0.5–4% anthraquinone; wt./wt.) during repellent exposure. Male turkeys exhibited 78–99% avoidance of food treated with 0.5–4% anthraquinone. Female and male turkeys that consumed more than 200 mg and 100 mg of anthraquinone, respectively, subsequently avoided food treated only with an UV-absorbent cue. In contrast, unconditioned females consumed 58% more food treated with the UV-absorbent cue than untreated food. Thus, wild turkeys do not prefer foods associated with UV wavelengths regardless of feeding experience. We also observed 1) a weak negative correlation between body condition and intestinal parasite infection and 2) moderate, positive correlations between consumption of food treated with the conditioned UV cue and intestinal parasite infection among male turkeys. The UV feeding cue was used to maintain food avoidance during the four days subsequent to postingestive conditioning. Moreover, the consequences of consuming food treated with the postingestive, UV-absorbent repellent were necessary for conditioned avoidance of the UV-absorbent cue. These findings suggest functional significance of UV feeding cues for avian foraging behavior, the implications of which will enable subsequent investigations regarding the sensory physiology and behavioral ecology of wild birds.

Published by Elsevier Inc.

## 1. Introduction

Most birds appear to be capable of sensing UV visual signals [1], but little is known about how they functionally use this information,

\* Corresponding author. Tel.: +1 970 266 6136; fax: +1 970 266 6138.

E-mail address: [Scott.J.Werner@aphis.usda.gov](mailto:Scott.J.Werner@aphis.usda.gov) (S.J. Werner).

particularly in the context of foraging. Ultraviolet cues could be used for foraging in two ways: 1) to detect foraging patches and recognize individual food items, and 2) to assess the relative quality of food items [2]. Comparative studies have found that not all bird species that could benefit from the use of UV feeding cues have evolved the retinal color receptors to do so (e.g. plunge-diving seabirds; [3]). Intraspecific studies have demonstrated that some bird species do indeed use UV cues to detect their food. Diurnal, predatory birds such as the Eurasian kestrel (*Falco tinnunculus*), rough-legged buzzard (*Buteo lagopus*) [4] and the great grey shrike (*Lanius excubitor*; [5]) use the UV reflectance of rodent urine to choose foraging patches where they are more likely to find these prey. Similarly, blue tits (*Parus caeruleus*) are able to find the first of a set of experimentally hidden cabbage moth (*Mamestra brassicae*) caterpillars more quickly with UV illumination than without it [6].

Many of the fruits eaten by birds exhibit high UV contrast with their backgrounds [7,8]. In a field study where UV filters were placed over *Psychotria emetica*, a tropical understory shrub, fewer fruits were taken when UV irradiance onto fruits was blocked compared to when UV transmitting filters were used [9]. Of course birds are not the only taxa to rely upon UV cues to detect their food. Predatory jumping spiders (*Portia labiata*) are preferentially attracted to the webs of their prey spider (*Argiope versicolor*), but only when the web reflects UV wavelengths [10]. Thus, birds and other animals can detect food more easily using UV cues. It is not clear, however, if birds use UV cues to assess the quality of their food.

Although both the strength of UV reflectance and predator preferences are often positively associated with specific prey, it is not known if preferences associated with UV reflectance increase the lifetime fitness of the forager. Are UV-reflecting prey more nutritious (*sensu lato*)? For example, are the prey biases observed among kestrels, for male rodents and for certain rodent species (see review; [2]), simply due to differences in signal detectability (i.e. greater UV reflectance) or have these predators learned that prey that exhibit greater UV reflectance provide greater benefits (e.g. more fat resources or fewer parasites)? Unfortunately very little is known about how birds utilize UV feeding cues; are there innate preferences for UV-reflecting or UV-absorbing food, or do birds learn to associate UV cues with food quality?

Ecologically-relevant, newborn color preferences and ontogenetic changes in color preferences have been studied experimentally in birds using only human-perceived colors (400–700 nm). Because of their experimental tractability, most of these studies have used domestic fowl (*Gallus gallus domesticus*) chicks as study subjects. Newborn domestic chicks prefer food items that are red or green in color if they are fruit-shaped, but avoid red items that are insect-shaped [11]. Chicks learn more easily to avoid distasteful food items that are red or yellow [12], or that contrast with their background [13], but some combinations of color and palatability are difficult for them to learn. For example, chicks require exposure to high quinine concentrations in their prey to learn that purple is unpalatable, but low quinine levels are sufficient for them to learn to avoid distasteful green prey [13].

Ontogenetic differences have been observed in UV foraging preference in redwings (*Turdus iliacus*; [14]). They discovered that wild-caught adult redwings preferred UV-reflecting bilberry (*Vaccinium myrtillus*) fruits over bilberries whose UV-reflecting waxy coat had been removed, but only when UV illumination was provided. Naïve, captive-reared redwing juveniles, however, showed no preference for the UV-reflecting fruits in either lighting regime, suggesting that redwings must learn to prefer UV wavelengths (or that their UV perception develops later in life). Ripe fruits often reflect more UV wavelengths [9], possibly explaining why many birds are attracted positively to UV wavelengths. Alternatively, plants may have co-opted existing avian preferences for UV-reflecting mates through sensory exploitation [15] in order to achieve greater seed dispersal by avian frugivores. Others posit that UV wavelengths have no special “meaning” via sensory bias [16], but are simply another color for which birds must learn context

dependency (just as birds must learn that some red fruits are unpalatable; [17]). To better understand how birds can use UV feeding cues, we experimentally investigated the foraging behavior of avian subjects with UV vision.

We used the wild turkey (*Meleagris gallopavo*) to investigate the functional significance of UV feeding cues. Wild turkeys are omnivores who consume a wide variety of vegetation, fruits, seeds, insects and other invertebrates [18]. Several lines of evidence support our contention that UV vision is important to turkey natural history. First, domestic turkeys (*M. gallopavo*) are attracted to housing with UV lighting [19]. Second, although they lack UV-sensitive opsin photopigments, ocular oil droplets associated with their short-wavelength sensitive cones apparently permit UV vision [20]. Domestic turkeys have considerable sensitivity to wavelengths in the UV-A spectral range (315–400 nm; [20]). Increment threshold psychophysiological tests have shown that domestic turkey poults are maximally sensitive to the UV spectrum at 380 nm [16]. Other studies have demonstrated that UV vision is probably of relevance to the social and sexual interactions of turkeys as well. The intensity of the UV reflectance of iridescent feathers from male wild turkeys is condition-dependent [21] and the plumage of domestic turkey poults exhibits UV-reflective patterning that is associated with body sites of harmful pecking in commercial poultry houses [22]. Moreover, another wild species in the order Galliformes, the black grouse (*Tetrao tetrix*), prefers UV-reflecting morphs of a fruit that is a seasonally important component of their diet [23].

Because the implications of UV cues are poorly understood for avian foraging behavior, we compared the feeding response of conditioned and unconditioned wild turkeys offered food treated with an UV-absorbent cue subsequent to conditioning with an UV-absorbent, postingestive repellent. If wild birds prefer foods associated with UV wavelengths regardless of feeding experience (hypothesis 1), then conditioned and unconditioned wild turkeys will prefer foods treated with an UV cue. If UV feeding cues, like other visual and gustatory cues [24,25], are used functionally for avian foraging behavior (hypothesis 2), then wild turkeys conditioned with an UV-absorbent, postingestive repellent will subsequently avoid food treated with an UV-absorbent cue, even in the absence of the aversive consequence.

Although intestinal parasite infection (e.g. *Eimeria* spp.) decreases food consumption in domestic turkeys [26–29], the effects of body condition and parasite load are poorly understood for the process of food selection. Coccidia infection influences sexual selection among female wild turkeys [30] and UV plumage signaling among male wild turkeys [21]. Body condition or parasite infection of wild turkeys may also influence an individual's selection of food treated with an UV cue previously paired with negative postingestive consequences. If body condition or parasite infection influences the process of avian food selection (hypothesis 3), then consumption of food treated with an aversively-conditioned UV cue will be least among wild turkeys with poor body condition or high parasite infection.

## 2. Feeding experiments

### 2.1. Subjects and testing facilities

Wild turkeys (4–6 years of age) were maintained at the Department of Biology's Avian Research Facility at the University of Mississippi Field Station in Lafayette County, Mississippi, USA. The wild turkey flock of game farm origin was raised in captivity from hatching. Twenty netted enclosures (4.0 × 3.7 × 1.8 m) were established within a 0.04-ha flight pen for the study of hens (i.e. female wild turkeys; body mass average = 4.07 kg, range = 3.02–5.75 kg). We used 16 individual cages (2.4 × 1.5 × 1.8 m) within an open-sided research aviary for the study of gobblers (male wild turkeys; body mass average = 9.87 kg, range = 7.45–11.50 kg). Clean water was provided ad libitum to all test subjects throughout the study.

## 2.2. Experimental procedures

Experimental investigation of foraging behavior requires that test subjects be exposed to ecologically relevant feeding conditions, but scientific ethics require that we minimize and mitigate the pain and distress of test subjects. In concert with the university veterinarian, we developed a protocol to meet both of these scientific needs. In the weeks prior to our study, all test subjects were offered a balanced poultry ration *ad libitum* to ensure that they were in the best condition for our study. We delayed our study until all test subjects had completed their molt. We paired hens within test cages to alleviate distress of individuals and disruption of flock dominance. We selected concentrations of test materials that had been previously approved for and tested with wild birds [31–33] to effectively condition and test avoidance whilst minimizing exposure among test subjects. The health of all subjects was monitored daily by study personnel and university animal care staff. Veterinary intervention due to our experimental procedures was never necessary. In accordance with U.S. federal law, all procedures were conducted only after review by and approval from the University of Mississippi's Institutional Animal Care and Use Committee (protocol #12-001; R. Buchholz – Study Director).

### 2.2.1. Exposure to an UV-absorbent, postingestive repellent

An anthraquinone-based repellent (Avipel®; Arkion Life Sciences, New Castle, DE, U.S.A.) was used to condition food avoidance among wild turkeys in captivity. Anthraquinone is a cathartic purgative [34] and is the active ingredient of avian repellents developed for the protection of rice [35–38], turf [39,40], corn [41] and sunflower crops [31,32]. We previously used spectrophotometry to determine that Avipel repellent absorbs UV wavelengths [33] throughout the spectrum visible to *M. gallopavo* (i.e. 315–400 nm; [20]).

Female wild turkeys ( $N = 40$ , experimentally naïve) acclimated within group cages (two hens per cage) and male turkeys ( $N = 16$ , experimentally naïve) acclimated within individual cages for five days prior to the study. During the acclimation period, one food bowl (1 kg untreated oats for hens, 0.5 kg untreated oats for gobblers) was presented in each cage at approximately 0800 h, daily.

Following acclimation, one bowl (1 kg untreated oats for hens, 0.5 kg untreated oats for gobblers) was offered in each cage at approximately 0800 h, daily for three days. Daily oat consumption was measured within each cage, including spillage, throughout the three-day pre-test. Paired hens and individual gobblers were ranked based upon average pre-test consumption and assigned to test groups (five groups of hens, four groups of gobblers) such that each group was similarly populated with turkeys that exhibited high–low daily consumption [31–33]. Test treatments were randomly assigned among groups.

On the day subsequent to the pre-test, one bowl (1 kg oats for hens, 0.5 kg oats for gobblers) was offered in each cage at approximately 0800 h. Turkeys in treatment groups one–four ( $n =$  four cages of paired hens per group;  $n =$  four individually-caged gobblers per group) received one bowl of oats treated with 0.5%, 1%, 2%, or 4% anthraquinone (wt./wt.) during the one-day test, respectively. We formulated oat treatments by applying aqueous suspensions (100 ml suspension/kg) to whole oats using a rotating mixer and household spray equipment [24,25,31,32]. Tested anthraquinone concentrations were based upon those previously used to develop anthraquinone concentration–response relationships for ring-necked pheasants (*Phasianus colchicus*), Canada geese (*Branta canadensis*), red-winged blackbirds (*Agelaius phoeniceus*) [31] and common grackles (*Quiscalus quiscula*) [32]. The availability of test subjects limited the control group (0% anthraquinone) to female turkeys; thus, the fifth group of hens ( $n =$  four control cages) again received 1 kg of untreated oats during the test. Daily oat consumption was measured within each cage, including spillage, at approximately 0800 h on the day subsequent to repellent exposure.

### 2.2.2. Conditioned avoidance of an UV-absorbent feeding cue

A titanium dioxide cue (Aeroxide® P25; Acros Organics, Fair Lawn, NJ, U.S.A.) was used to test food avoidance previously conditioned with the anthraquinone-based repellent. We previously used spectrophotometry to determine that this titanium dioxide cue absorbs UV wavelengths similarly to Avipel repellent [33] and throughout the spectrum visible to *M. gallopavo*.

Two bowls (1 kg oats per bowl for hens, 0.5 kg oats per bowl for gobblers) were offered in each cage at approximately 0800 h, daily, during the four days subsequent to repellent exposure. One bowl contained untreated oats. The alternate bowl contained oats treated only with the UV-absorbent cue (0.2% titanium dioxide, wt./wt.; [33]). We formulated oat treatments by applying aqueous suspensions (85 ml suspension/kg) to whole oats using a rotating mixer and household spray equipment. The east–west placement of treated and untreated oats was randomized on test day one, and was thereafter alternated daily, throughout the test. Daily consumption of treated and untreated oats was independently measured within each cage, including spillage, at approximately 0800 h throughout the test of conditioned avoidance.

## 2.3. Analytical chemistry

Reversed-phase, high performance liquid chromatography (HPLC) with UV detection (254 nm) was used to quantify anthraquinone residues for all repellent-treated oats ( $\pm 100$  ppm anthraquinone). We collected a 200 g sample of each treatment used for repellent exposure. Subsequent to formulations, all samples were transferred to a 4 °C refrigerator at the National Wildlife Research Center (Fort Collins, CO, U.S.A.) where they were stored for the duration of the analysis period.

Triplicate subsamples from each repellent treatment were extracted and analyzed. All samples were cryogenically homogenized. Control samples were fortified with 1,500 ppm and 40,000 ppm anthraquinone, and extracted to determine the recovery rate for the assay. We weighed 0.5 ( $\pm 0.05$ ) g of ground whole oats into 25-ml glass test tubes fitted with Teflon lined caps. We pipetted 8 ml of 25% hexane in chloroform (vol/vol) into each tube. Extraction was accomplished by vortexing each tube for 20 s, placing on a horizontal shaker for 30 min, sonicating for 30 min, and then centrifuging at 2,000 rpm for 10 min. The supernatant was carefully filtered through a 0.45  $\mu$ m Teflon filter into a 25-ml volumetric flask. The entire extraction procedure was replicated three times and the supernatants were combined. The sample was diluted to volume with the 25% hexane in chloroform solution and an aliquot was placed in a clean 25-ml glass test tube. The aliquot was evaporated to dryness at 50 °C under a gentle stream of nitrogen. The extract was reconstituted using 10 ml of methanol, sonicated for 30 min, and again centrifuged at 2,000 rpm for 10 min. Sample solutions were transferred into autosampler vials and analyzed by HPLC using an Agilent 1200 liquid chromatograph (Agilent Technologies, Inc., Santa Clara, CA, U.S.A.).

The HPLC instrument included a Waters X-Bridge Phenyl column (2.5  $\mu$ m, 2.1  $\times$  50 mm). The mobile phase gradient included 90% Millipore water and 10% methanol at 0 and 2 min, 20% Millipore water and 80% methanol at 4 and 7 min, and 100% methanol at 10 min. The HPLC flow rate, injection volume, and temperature were 0.3 ml/min, 5  $\mu$ l and 40 °C, respectively. A four-point external calibration curve was used to calibrate our HPLC instrument. Samples were run in triplicate each day and we checked single calibration points upon each ten injections. The average response was plotted against anthraquinone concentrations. Linear regression was used to calculate anthraquinone concentrations among samples.

## 2.4. Statistical analyses

The dependent measure for the repellent exposure phase of our study was calculated as test consumption of anthraquinone-treated oats relative to average pre-test consumption of untreated oats (i.e. percent

repellency =  $(1 - (\text{test consumption} \times \text{pre-test consumption}^{-1})) \times 100$ ; [31,32]). Logarithmic regression procedures (SAS v9.2) were used to analyze repellency as a function of actual anthraquinone concentration ( $\pm 100$  ppm) and predict a threshold anthraquinone concentration (i.e. 80% repellency; [31,32]) for hens and gobblers. Descriptive statistics ( $\bar{x} \pm \text{SE}$ ) were used to summarize oat and anthraquinone consumption during repellent exposure.

The dependent measure for the test of conditioned avoidance was average daily consumption of untreated oats and oats treated with the UV-absorbent cue throughout the test. Test consumption data for hens and gobblers were subjected to a repeated measures ANOVA. The random effect of our model was cages (i.e. paired hens, individual gobblers), the between-subjects effects were oat treatments (treated, untreated) and test groups (i.e. previous exposure to 0%, 0.5%, 1%, 2%, or 4% anthraquinone-treated oats), and the within-subject effect was test day. The group-by-treatment interaction was analyzed using the mixed procedure (SAS v9.2). Tukey's tests were used to separate the means of significant interactions ( $\alpha = 0.05$ ), and descriptive statistics ( $\bar{x} \pm \text{SE}$ ) and preference ratios [daily average  $\text{TiO}_2$  consumption  $\times$  (daily average  $\text{TiO}_2$  consumption + untreated consumption) $^{-1}$ ] were used to summarize and illustrate test consumption, respectively.

To test our prediction regarding the influence of subject body condition, we measured the condition (body mass  $\times$  tarsus length $^{-1}$ ) and enumerated intestinal parasites (*Eimeria* spp., *Capillaria* spp., other nematodes) from collected fecal samples [30] for each tested gobbler (i.e. independent of test groups;  $n = 16$ ). Body condition and parasite data were not available for individual hens that were paired for our feeding experiments. These indices of body condition were correlated with 4-day average test consumption and relative test consumption [4-day average  $\text{TiO}_2$  consumption  $\times$  (4-day average  $\text{TiO}_2$  consumption + untreated consumption) $^{-1}$ ] of food treated with the UV cue during the test of conditioned avoidance.

### 3. Results

#### 3.1. Exposure to an UV-absorbent, postingestive repellent

Hens in the control group consumed  $216.1 \pm 21.0$  g of untreated oats during the exposure phase of our study; their average, pre-test consumption of untreated oats was  $193.3 \pm 22.5$  g. In contrast, hens exposed to oats treated with 0.5–4% anthraquinone exhibited 75–98% repellency during repellent exposure (Table 1). Hen repellency ( $y$ ) was a function of anthraquinone concentration ( $x$ ):  $y = 10.746 \ln(x) - 12.029$  ( $r^2 = 0.94$ ,  $P = 0.030$ ). We therefore predicted a threshold concentration of 5,300 ppm anthraquinone (i.e. 80% repellency), or  $47.0 \pm 18.3$  mg anthraquinone  $\times$  kg body mass $^{-1}$ , for hens offered treated oats.


Gobblers exposed to oats treated with 0.5–4% anthraquinone exhibited 78–99% repellency during repellent exposure (Table 1). Gobbler

repellency ( $y$ ) was a function of anthraquinone concentration ( $x$ ):  $y = 9.921 \ln(x) - 2.260$  ( $r^2 = 0.93$ ,  $P = 0.034$ ). We therefore predicted a threshold concentration of 4000 ppm anthraquinone, or  $13.7 \pm 8.3$  mg anthraquinone  $\times$  kg body mass $^{-1}$ , for gobblers offered treated oats.

On average, hens and gobblers consumed  $114 \pm 88$  mg and  $48 \pm 15$  mg of anthraquinone when exposed to oats treated with 4% anthraquinone, respectively. In comparison, average consumption among hens and gobblers was  $204 \pm 34$  mg and  $129 \pm 38$  mg anthraquinone, respectively, when exposed to oats treated with 0.5%, 1%, or 2% anthraquinone. Thus, conditioned food avoidance was positively related to the amount of the postingestive repellent consumed during the one-day exposure.

#### 3.2. Conditioned avoidance of an UV-absorbent feeding cue

The five test groups of hens consumed different amounts of oats treated with the UV-absorbent cue and untreated oats during the four-day test of conditioned avoidance ( $F_{9,27} = 11.66$ ,  $P < 0.0001$ ; Fig. 1a). Unconditioned (control) hens consumed similar amounts of untreated



**Fig. 1.** Preference ratios for (a) five test groups of female wild turkeys (*Meleagris gallopavo*;  $n =$  four cages of paired hens per group) and (b) four test groups of male wild turkeys ( $n =$  four individually-caged gobblers per group) offered untreated food and food treated with an UV-absorbent cue (a.i. titanium dioxide; Acros Organics, Fair Lawn, NJ, U.S.A.) subsequent to one-day exposure to an UV-absorbent, postingestive repellent (a.i. 9,10-anthraquinone; Arkion Life Sciences, New Castle, DE, U.S.A.). Preference ratios  $>0.5$  indicate preference for treated food; ratios  $<0.5$  indicate avoidance of treated food. The control group of females (unfilled squares) was exposed to untreated oats and test groups (filled shapes) were exposed to 0.5%, 1%, 2%, or 4% anthraquinone (AQ, wt./wt.) prior to the test of conditioned avoidance.

**Table 1**

Feeding repellency of oats treated with an anthraquinone-based repellent (Avipel®; Arkion Life Sciences, New Castle, DE, U.S.A.) among wild turkeys, *Meleagris gallopavo*. Actual anthraquinone concentrations among oat seed treatments were quantified using high performance liquid chromatography. The method detection limit (MDL) of our analyses was 0.50  $\mu\text{g}$  anthraquinone/g. Percent repellency represents daily consumption of repellent-treated oats relative to average pre-treatment consumption of untreated oats among five groups of females ( $n =$  four cages of paired hens per group) and four groups of males ( $n =$  four individually-caged gobblers per group).

Targeted anthraquinone concentration (%)	Actual anthraquinone concentration (ppm)	Hen repellency (%)	Gobbler repellency (%)
0	<MDL	12	
0.5	4100	75	78
1	8800	89	91
2	19,100	95	97
4	34,400	98	99



oats and oats treated with titanium dioxide throughout the test (Fig. 1a). The control group consumed an average of  $138.6 \pm 13.1$  g of oats treated with titanium dioxide and  $87.7 \pm 17.5$  g of untreated oats per day (Tukey  $P = 0.667$ ). Thus, unconditioned wild turkeys did not significantly prefer foods treated with an UV feeding cue.

In contrast, hens conditioned with the UV-absorbent, postingestive repellent subsequently avoided oats treated with the UV-absorbent cue throughout the test (Fig. 1a). Hens previously exposed to oats treated with 0.5% anthraquinone consumed an average of  $51.0 \pm 17.1$  g of oats treated with titanium dioxide and  $197.6 \pm 21.5$  g of untreated oats per day (Tukey  $P < 0.001$ ). The group of hens exposed to oats treated with 1% anthraquinone subsequently consumed an average of  $34.8 \pm 17.9$  g of oats treated with titanium dioxide and  $229.7 \pm 26.4$  g of untreated oats per day (Tukey  $P < 0.0001$ ). Hens previously exposed to oats treated with 2% anthraquinone consumed an average of  $78.3 \pm 16.5$  g of oats treated with titanium dioxide and  $173.7 \pm 24.7$  g of untreated oats per day (Tukey  $P = 0.038$ ). Thus, the UV-absorbent cue was used to maintain avoidance during the four days subsequent to postingestive conditioning.

The group of hens previously exposed to oats treated with 4% anthraquinone consumed an average of  $81.9 \pm 13.7$  g of oats treated with titanium dioxide and  $128.1 \pm 14.9$  g of untreated oats per day (Tukey  $P = 0.774$ ). Thus, conditioned avoidance of food treated with the UV-absorbent cue was influenced by the amount of repellent-treated oats consumed during exposure (i.e. the negative postingestive consequence).

The four test groups of gobblers also consumed different amounts of oats treated with the UV-absorbent cue and untreated oats during the four-day test of conditioned avoidance ( $F_{7,21} = 14.20$ ,  $P < 0.0001$ ; Fig. 1b). Gobblers previously exposed to oats treated with 0.5% anthraquinone consumed an average of  $4.2 \pm 4.3$  g of oats treated with titanium dioxide and  $196.0 \pm 18.6$  g of untreated oats per day (Tukey  $P < 0.0001$ ). The group of gobblers exposed to oats treated with 1% anthraquinone subsequently consumed an average of  $86.6 \pm 23.6$  g of oats treated with titanium dioxide and  $117.4 \pm 28.0$  g of untreated oats per day (Tukey  $P = 0.961$ ). Two gobblers in the group previously exposed to 1% anthraquinone consumed more oats treated with the UV-absorbent cue than untreated oats on test days one–four, and test days one, two and four, respectively. Of these two gobblers, one had the highest parasite infection measured in the study [i.e. greatest abundance of *Eimeria* spp. (203/fecal g), *Capillaria* spp. (1267/g) and other nematodes (34/g)] and the other gobbler had an intermediate parasite infection among tested gobblers.

Gobblers previously exposed to oats treated with 2% anthraquinone consumed an average of  $12.4 \pm 7.6$  g of oats treated with titanium dioxide and  $206.3 \pm 24.5$  g of untreated oats per day (Tukey  $P < 0.0001$ ). The group of gobblers exposed to oats treated with 4% anthraquinone subsequently consumed an average of  $77.4 \pm 21.6$  g of oats treated with titanium dioxide and  $175.3 \pm 25.8$  g of untreated oats per day (Tukey  $P = 0.054$ ). Two gobblers in the group previously exposed to 4% anthraquinone consumed more oats treated with the UV-absorbent cue than untreated oats on test days two and four, and test days two, three and four, respectively. Of these two gobblers, one consumed the least amount (0.7 g) of 4% anthraquinone-treated oats during repellent exposure. Similar to the hens, conditioned avoidance of UV-absorbent food was influenced by the amount of repellent-treated oats consumed by tested gobblers during exposure.

With further regard to the relationship between body condition and conditioned avoidance of food treated with an UV cue, we observed moderate, positive correlations [42] between consumption of food treated with the conditioned UV cue and intestinal parasite infection among tested gobblers (Table 2). We also observed a weak negative correlation between body condition and intestinal parasite infection (Table 2). Thus, intestinal parasites moderately decreased conditioned avoidance of food treated with an UV cue previously paired with negative postingestive consequences during the gobbler test.

**Table 2**

Correlation coefficients for empirical relationships between body condition (body mass  $\times$  tarsus length<sup>-1</sup>), intestinal parasite infection (abundance  $\times$  fecal g<sup>-1</sup>), and test consumption and relative test consumption of food treated with an UV cue among male wild turkeys, *Meleagris gallopavo*, used to test conditioned avoidance of food treated with an UV cue previously associated with negative postingestive consequences.

	Body condition	<i>Eimeria</i> spp.	<i>Capillaria</i> spp.	Other nematodes
Consumption of UV-treated food	0.013	0.496	0.479	0.433
Relative consumption of UV-treated food	−0.002	0.519	0.503	0.452
<i>Eimeria</i> spp.	−0.213			
<i>Capillaria</i> spp.	−0.219			
Other nematodes	−0.249			

#### 4. Discussion

Female turkeys exhibited no avoidance of untreated food and 75–98% avoidance of food treated with an UV-absorbent, postingestive repellent (0.5–4% anthraquinone; wt./wt.) during one day of repellent exposure. Male turkeys exhibited 78–99% avoidance of food treated with 0.5–4% anthraquinone. Hens and gobblers that consumed more than 200 mg and 100 mg of the UV-absorbent, postingestive repellent, respectively, subsequently avoided food treated only with an UV-absorbent cue. Ultraviolet feeding cues were therefore specifically related to the postingestive consequences of the subsequent reinforcer [43]. In contrast, unconditioned hens consumed 58% more food treated with the UV-absorbent cue than untreated food. Thus, conditioned food avoidance was positively related to the amount of the postingestive repellent consumed during the one-day exposure, and the consequences of consuming oats treated with the postingestive, UV-absorbent repellent were necessary for conditioned avoidance of the UV-absorbent cue. Wild turkeys do not prefer foods associated with UV wavelengths regardless of feeding experience (hypothesis 1).

In the absence of negative postingestive feedback [44,45], UV feeding cues are therefore unlikely to function as aposematic signals [46] or elicit food avoidance in wild birds. Ultraviolet foraging behavior is therefore a function of its consequences [47] and UV feeding cues are used functionally for foraging behavior in wild turkeys (hypothesis 2). Subsequent investigations should be focused to relate food preference with the chromatic and achromatic characteristics of natural foods [7]. Newborn and ontogenetic color preferences can be better understood by investigating the full spectrum visible to and used by avian subjects.

We predicted that consumption of food treated with an aversively-conditioned UV cue would be least among wild turkeys with poor body condition or high parasite infection (Hypothesis 3). Rather, we observed moderate, positive correlations between consumption of food treated with the conditioned UV cue and intestinal parasite infection (Table 2). In context of food selection, aversive feedback or a lack of positive feedback from the gut to the central nervous system causes animals to reduce food consumption [44,45]. Perhaps poor body condition or high parasite infection can interfere with feedback-mediated consumption of foods, including those previously associated with negative postingestive consequences. Supplemental studies are recommended to further investigate the influence of parasite infection and subject body condition for the process of avian food selection.

In conclusion, we discovered that wild turkeys do not prefer foods associated with UV wavelengths regardless of feeding experience. Rather, we found that wild turkeys can use UV feeding cues to avoid foods previously associated with negative postingestive consequences, and that this cue–consequence association was dependent upon the amount of previously experienced, postingestive consequences. Thus, UV feeding cues, like other visual and gustatory cues, have functional significance for avian foraging behavior. Not all individuals in our study, however, exhibited conditioned avoidance of foods treated with an UV feeding cue, an effect moderately related to intestinal parasite infection. Our study of the functional use of UV feeding cues in wild

turkeys contributes to a broader avian data set and will enable subsequent investigations regarding the sensory physiology and behavioral ecology of wild birds.

## Acknowledgments

We thank M. L. Avery, B. F. Blackwell, B. A. Kimball, G. M. Linz, F. D. Provenza, and M. E. Tobin for their thoughtful review of our manuscript. P. B. Fioranelli (Mississippi Field Station — National Wildlife Research Center, Starkville, MS, U.S.A.) provided dedicated assistance with constructing and disassembling enclosures for our hen study. We also thank the analytical chemistry unit at the National Wildlife Research Center (Fort Collins, CO, U.S.A.) for performing anthraquinone residue analyses. Arkion Life Sciences (New Castle, DE, U.S.A.) provided the Avipel® repellent and Acros Organics (Fair Lawn, NJ, U.S.A.) provided the Aeroxide® P25 for our study. Corporate collaborations do not imply endorsement by the United States Department of Agriculture. This project was supported in part by California Department of Food and Agriculture's Specialty Crop Block Grant to S.J.W. (Grant Agreement SCB10034).

## References

- [1] Aidala Z, Huynen L, Brennan PLR, Musser J, Fidler A, Chong N, et al. Ultraviolet visual sensitivity in three avian lineages: paleognaths, parrots, and passerines. *J Comp Physiol A* 2012;198:495–510.
- [2] Honkavaara J, Koivula M, Korpimäki E, Siitari H, Viitala J. Ultraviolet vision and foraging in terrestrial vertebrates. *Oikos* 2002;98:505–11.
- [3] Hästad O, Ernstdotter E, Odeen A. Ultraviolet vision and foraging in dip and plunge diving birds. *Biol Lett* 2005;1:306–9.
- [4] Viitala J, Korpimäki E, Palokangas P, Koivula M. Attraction of kestrels to vole scent marks visible in ultraviolet light. *Nature* 1995;373:425–7.
- [5] Probst R, Pavlicev M, Viitala J. UV reflecting vole scent marks attract a passerine, the great grey shrike (*Lanius excubitor*). *J Avian Biol* 2002;33:437–40.
- [6] Church SC, Bennett ATD, Cuthill IC, Partridge JC. Ultraviolet cues affect the foraging behaviour of blue tits. *Proc R Soc Lond B* 1998;265:1509–14.
- [7] Schaefer HM, Levey DJ, Schaefer V, Avery ML. The role of chromatic and achromatic signals for fruit detection by birds. *Behav Ecol* 2006;17:784–9.
- [8] Schaefer HM, McGraw K, Catoni C. Birds use fruit colour as honest signal of dietary antioxidant rewards. *Funct Ecol* 2008;22:303–10.
- [9] Altshuler DL. Ultraviolet reflectance in fruits, ambient light composition and fruit removal in a tropical forest. *Evol Ecol Res* 2001;3:767–78.
- [10] Zou Y, Araujo DP, Lim MLM, Li D. Ultraviolet is a more important cue than reflection in other wavelengths for a jumping spider to locate its spider prey. *Anim Behav* 2011;82:1457–63.
- [11] Gamberale-Stille G, Tullberg BS. Fruit or aposematic insect? Context-dependent colour preferences in domestic chicks. *Proc Biol Sci* 2001;268:2525–9.
- [12] Rowe C, Skelhorn J. Colour biases are a question of taste. *Anim Behav* 2005;69:587–94.
- [13] Halpin CG, Skelhorn J, Rowe C. Being conspicuous and defended: selective benefits for the individual. *Behav Ecol* 2008;19:1012–7.
- [14] Siitari H, Honkavaara J, Viitala J. Ultraviolet reflection of berries attracts foraging birds. A laboratory study with redwings (*Turdus iliacus*) and bilberries (*Vaccinium myrtillus*). *Proc R Soc Lond* 1999;266:2125–9.
- [15] Bennet ATD, Théry M. Avian color vision and coloration: multidisciplinary evolutionary biology. *Am Nat* 2007;169:S1–6.
- [16] Barber CL, Prescott NB, Jarvis JR, Le Sueur C, Perry GC, Wathes CM. Comparative study of the photopic spectral sensitivity of domestic ducks (*Anas platyrhynchos domesticus*), turkeys (*Meleagris gallopavo gallopavo*) and humans. *Br Poultry Sci* 2006;47:365–74.
- [17] Lev-Yadun S, Ne'eman G, Izhaki I. Unripe red fruits may be aposematic. *Plant Signal Behav* 2009;4:836–41.
- [18] Hurst GA. Foods and feeding. In: Dickson JG, editor. *The wild turkey: biology and management*. Harrisburg, PA: Stackpole Books; 1992. p. 66–83.
- [19] Moinard C, Sherwin CM. Turkeys prefer fluorescent light with supplementary ultraviolet radiation. *Appl Anim Behav Sci* 1999;64:261–7.
- [20] Hart NS, Partridge JC, Cuthill IC. Visual pigments, cone oil droplets, ocular media and predicted spectral sensitivity in the domestic turkey (*Meleagris gallopavo*). *Vision Res* 1999;39:3321–8.
- [21] Hill GE, Doucet SM, Buchholz R. The effect of coccidial infection on iridescent plumage coloration in wild turkeys. *Anim Behav* 2005;69:387–94.
- [22] Sherwin CM, Devereux CL. A preliminary investigation of ultraviolet-visible markings in domesticated turkey chicks and a possible role in injurious pecking. *Br Poultry Sci* 1999;40:429–33.
- [23] Siitari H, Viitala J, Hovi M. Behavioural evidence for ultraviolet vision in a tetraonid species — foraging experiment with black grouse (*Tetrao tetrix*). *J Avian Biol* 1999;30:199–202.
- [24] Werner SJ, Kimball BA, Provenza FD. Food color, flavor, and conditioned avoidance among red-winged blackbirds. *Physiol Behav* 2008;93:110–7.
- [25] Werner SJ, Provenza FD. Reconciling sensory cues and varied consequences of avian repellents. *Physiol Behav* 2011;102:158–63.
- [26] Clarkson MJ. The life history and pathogenicity of *Eimeria meleagridis* Tyzzer 1929, in the turkey poult. *Parasitology* 1959;49:70–82.
- [27] Hein H. *Eimeria adenoeides* and *E. meleagridis*: pathogenic effect in turkey poults. *Exp Parasitol* 1969;24:163–70.
- [28] Yvone P, Berdougou H, Naciri M, Bree A, Lafont JP. Pathogenic study of turkey coccidiosis due to *Eimeria adenoeides*. *Ann Rech Vet* 1978;9:531–9.
- [29] Augustine PC, Thomas OP. *Eimeria meleagridis* in young turkeys: effects on weight, blood, and organ parameters. *Avian Dis* 1979;23:854–62.
- [30] Buchholz R. Effects of parasitic infection on mate sampling by female wild turkeys (*Meleagris gallopavo*): should infected females be more or less choosy? *Behav Ecol* 2004;15:687–94.
- [31] Werner SJ, Carlson JC, Tupper SK, Santer MM, Linz GM. Threshold concentrations of an anthraquinone-based repellent for Canada geese, red-winged blackbirds, and ring-necked pheasants. *Appl Anim Behav Sci* 2009;121:190–6.
- [32] Werner SJ, Linz GM, Carlson JC, Pettit SE, Tupper SK, Santer MM. Anthraquinone-based bird repellent for sunflower crops. *Appl Anim Behav Sci* 2011;129:162–9.
- [33] Werner SJ, Tupper SK, Carlson JC, Pettit SE, Ellis JW, Linz GM. The role of a generalized ultraviolet cue for blackbird food selection. *Physiol Behav* 2012;106:597–601.
- [34] Fraser CM, editor. *The Merck veterinary manual*. 7th ed. Rahway, NJ: Merck & Company, Inc.; 1991.
- [35] Avery ML, Humphrey JS, Decker DG. Feeding deterrence of anthraquinone, anthracene, and anthrone to rice-eating birds. *J Wildl Manag* 1997;61:1359–65.
- [36] Avery ML, Humphrey JS, Primus TM, Decker DG, McGrane AP. Anthraquinone protects rice seed from birds. *Crop Prot* 1998;17:225–30.
- [37] Cummings JL, Avery ML, Mathre O, Wilson EA, York DL, Engeman RM, et al. Field evaluation of Flight Control™ to reduce blackbird damage to newly planted rice. *Wildl Soc Bull* 2002;30:816–20.
- [38] Cummings JL, Byrd RW, Eddleman WR, Engeman RM, Tupper SK. Effectiveness of AV-1011® to reduce damage to drill-planted rice from blackbirds. *J Wildl Manag* 2011;75:353–6.
- [39] Dolbeer RA, Seamans TW, Blackwell BF, Belant JL. Anthraquinone formulation (Flight Control™) shows promise as an avian feeding repellent. *J Wildl Manag* 1998;62:1558–64.
- [40] Blackwell BF, Seamans TW, Dolbeer RA. Plant growth regulator (Stronghold™) enhances repellency of anthraquinone formulation (Flight Control™) to Canada geese. *J Wildl Manag* 1999;63:1336–43.
- [41] Blackwell BF, Helon DA, Dolbeer RA. Repelling sandhill cranes from corn: whole-kernel experiments with captive birds. *Crop Prot* 2001;20:65–8.
- [42] Cohen J. Statistical power analysis for the behavioral sciences. 2nd ed. Hillsdale, NJ: Lawrence Erlbaum Associates, Publishers; 1988.
- [43] Garcia J, Koelling RA. Relation of cue to consequence in avoidance learning. *Psychon Sci* 1966;4:123–4.
- [44] Provenza FD. Postingestive feedback as an elementary determinant of food preference and intake in ruminants. *J Range Manage* 1995;48:2–17.
- [45] Provenza FD, Villalba JJ. Foraging in domestic vertebrates: linking the internal and external milieu. In: Bels VL, editor. *Feeding in domestic vertebrates: from structure to function*. Oxfordshire: CABI; 2006. p. 210–40.
- [46] Lyytinen A, Alatalo RV, Lindstrom L, Mappes J. Can ultraviolet cues function as aposematic signals? *Behav Ecol* 2001;12:65–70.
- [47] Skinner BF. Selection by consequences. *Science* 1981;213:501–4.